

# 87736-344322-1-CE\_Turn.docx

*by* Sugiharto Totok

---

**Submission date:** 14-Mar-2024 02:39PM (UTC+0800)

**Submission ID:** 2209840614

**File name:** 87736-344322-1-CE\_Turn.docx (681.79K)

**Word count:** 5773

**Character count:** 34441

## **Methyl Jasmonate Elicitation Enhanced Biomass, Phenolic and Flavonoid Production of Adventitious Root Culture of *Gynura procumbens* in a Balloon Type Bubble Bioreactor**

Ely Tri Wijayanti<sup>1</sup>, Dannis Yuda Kusuma<sup>1</sup>, Alfinda Novi Kristanti<sup>2</sup>, Anjar Tri Wibowo<sup>1</sup>, Sugiharto<sup>1</sup>, Djarot Sugiarto<sup>3</sup>, Yosephine Sri Wulan Manuhara<sup>1\*</sup>

<sup>1</sup>Department of Biology, Faculty of Science and Technology, Universitas Airlangga, Surabaya, East Java, Indonesia

<sup>2</sup>Department of Analytic, Faculty of Science and Analytical Data, Institut Teknologi Sepuluh Nopember, Surabaya, East Java, Indonesia

<sup>3</sup>Department of Chemistry, Faculty of Science and Technology, Universitas Airlangga, Surabaya, East Java, Indonesia

\*yosephine-s-w-m@fst.unair.ac.id

### **ABSTRACT**

Secondary metabolites in the medicinal plant *Gynura procumbens* have various beneficial pharmacological activities. However, the limited availability of these beneficial compounds in nature has constrained the possibilities for therapeutic applications. The use of exogenously applied methyl jasmonate (MeJa) as an elicitor has been reported to trigger the accumulation of secondary metabolites in plant tissues cultured *in vitro*. This study aimed to assess how varying periods of MeJa elicitation affect the growth and accumulation of phenolic and flavonoid compounds in adventitious root cultures of *G. procumbens* within a balloon-type bubble bioreactor (BTBB). Adventitious roots were induced from leaf and stem explants obtained from established *in vitro* shoots. Upon transfer to the BTBB, the cultures were supplemented with 100  $\mu$ M MeJa and sampled at intervals of 1, 2, 3, and 4 weeks. The results revealed that the inclusion of MeJa in the culture medium significantly boosted both growth and the accumulation of phenolic and flavonoid compounds in *G. procumbens*. The highest biomass and concentrations of phenolic and flavonoid compounds were observed after 3 weeks of elicitation. These findings highlight the potential of MeJa elicitation for enhancing the production of valuable bioactive compounds in *G. procumbens* adventitious root cultures, showcasing its utility for industrial applications.

**Keywords:** balloon-type bubble bioreactor; flavonoid; *Gynura procumbens*; methyl jasmonate; phenol

## INTRODUCTION

*Gynura procumbens* is an indigenous Indonesian plant that has been traditionally used as a vegetable, anesthesia drug, and herbal medicine to treat various symptoms, such as fever, high cholesterol level, dysentery, tumors, and kidney problems. Previous works have reported the potential use of *G. procumbens* for cancer (Nurulita et al., 2012) and diabetic treatments (Algariri et al., 2013). It has also been widely used as an immunomodulator agent (Takanashi et al., 2019), vasodilatation agent (Hoe et al., 2011; Ng et al., 2013), antiviral agent (Jarikasem et al., 2013), anti-inflammatory agent (Iskander et al., 2002) and antioxidant agent (Krishnan et al., 2015). These beneficial pharmacological properties are mainly because of the flavonoid compounds present in the leaf and root organs of *G. procumbens* (Tan et al., 2016). These compounds include kaempferol, quercetin (Kim et al., 2011), and flavanon (Manogaran et al., 2019). Despite the potential of these compounds, their limited availability in nature has constrained the possibilities of these compounds for therapeutic applications.

*G. procumbens* conventionally propagated by cuttings. This method is slow and time-consuming since the growth of plants can be affected by season and environment. To meet the growing demand for bioactive compounds sourced from plants, the development of an efficient in vitro system using tissue culture techniques is crucial. Numerous plant secondary metabolites have been successfully generated through *in vitro* systems (Cui et al., 2010a), such as caffeic acid, anthraquinones phenolics, and flavonoids. Numerous secondary metabolites are biosynthesized by specialized cells. For example, adventitious root cultures have demonstrated the ability to yield a greater quantity of secondary metabolites compared to those obtained from cell cultures. Adventitious roots are frequently employed at an industrial scale for the production of these safe and non-carcinogenic secondary metabolites (Baque et al., 2014; 2013; Murthy et al., 2016). For instance, the FDA has approved the utilization of bioactive compounds derived from adventitious roots of *Panax ginseng* and *Echinacea purpurea* in the pharmaceutical and cosmetic industries. (Baque et al., 2014).

Adventitious roots from other plant species, such as *Hypricum perforatum* (Cui et al., 2010a), *Astragalus membranaceus* (Wu et al., 2011), and *G. procumbens* (Saiman et al., 2012), have also been reported.

*In vitro* plant culture systems employing bioreactors are commonly employed to augment the production of secondary metabolites. A bioreactor is an engineered system with the capability to sustain plant growth within varying chemical conditions, utilizing liquid culture media, and control physical factors such as airflow, aeration, temperature, pH, and dissolved oxygen levels. (Karuppusamy et al., 2009). (Karuppusamy, 2009). Different types of bioreactors have been employed in plant research. Of these, balloon-type bubble bioreactor (BTBB) is frequently used for adventitious root cultures (Baque et al., 2014; Choi et al., 2000; Lee et al., 2014; Lee & Paek, 2012; Paek et al., 2001). In our previous works, we have enhanced the biomass of *G. procumbens* adventitious root cultures in BTBB by 13-fold through the addition of 50% sucrose into the media (Manuhara et al., 2017; Muthoharoh et al., 2019). Besides sucrose, various biotic elicitors (microbial extract, polysaccharide, and oligosaccharide) and abiotic elicitors (heavy metal, pesticide, UV light, cold shock, hyperosmotic stress, and ultrasound) have been used to enhance the production of secondary metabolites. For example, heavy metal (CuSO<sub>4</sub>) and low concentration of *Saccharomyces cerevisiae* extract have been added to *G. procumbens* cultures to improve biomass and flavonoid production (Faizah et al., 2018). The addition of jasmonic acid (JA) or its methyl ester variant methyl jasmonate (MeJa) into *in vitro* plant cultures has been shown to effectively increase the secondary metabolite accumulation (Foyer & Harbinson, 2019; Ho et al., 2020; Rahimi et al., 2015). This promotive effect of JA or MeJa treatment has been shown in adventitious root cultures of *Panax ginseng* (Ali et al., 2007; Kim et al., 2004), *Pseudostellaria heterophylla* (Wang et al., 2015), *Eleutherococcus koreanum* (Lee et al., 2015) and *Oplonanax elatus* (Jiang et al., 2017).

In this study, we assessed how varying periods of MeJa elicitation affect the growth, as well as the production of phenolic and flavonoid compounds in adventitious roots of *G. procumbens* within a BTBB. Additionally, we examined changes in malondialdehyde (MDA) and proline levels in MeJa-treated adventitious roots, which serve as indicators of cellular oxidative stress. Furthermore, we analyzed alterations in pH, sugar levels, and

electrical conductivity of the culture media, providing valuable insights into the overall biochemical and environmental dynamics of the system.

## MATERIALS AND METHODS

### Materials

Leaf and stem tissues of *Gynura procumbens* (Lour.) Merr obtained from Purwodadi Botanical Garden, Pasuruan, East Java, Indonesia (Botanical Garden identification number/IPH.06/HM/XI/2015) were used to establish adventitious root cultures. Murashige and Skoog's solid medium were used to induce adventitious roots with 7 g/L agar.

### Methods

#### Induction of adventitious roots

The leaves and stems of *G. procumbens* were first cleaned with detergent for 5 minutes and then thoroughly rinsed with tap water. Subsequently, the cleaned leaves were immersed in a 10% (v/v) Clorox solution for 10 minutes (commercial bleach with 5.25% w/v sodium hypochlorite), followed by three washes with sterile distilled water. The plant explants were then sectioned into 1 cm<sup>2</sup> pieces and placed on Murashige and Skoog (MS) solid medium supplemented with 5 mg/L indole-3-butyric acid (IBA), 30 g/L sucrose, and 7 g/L agar for leaf-derived explants, while stem-derived explants were cultured on MS solid medium supplemented with 2 mg/L indole acetic acid (IAA), 4 mg/L 6-benzylaminopurine (BAP), 30 g/L sucrose, and 7 g/L agar. The pH of the medium was adjusted to 5.8 and autoclaved at 121°C for 15 minutes. The cultures were then incubated in darkness at 25±2°C. After 21 days of culture, the adventitious roots were collected for subsequent BTBB culture.

#### Adventitious roots culture in BTBB

Approximately 2 grams of adventitious roots were placed in a BTBB with a maximum capacity of 1 liter, containing 500 mL of liquid MS medium supplemented with 5 mg/L IBA and 30 g/L sucrose. The cultures were maintained in darkness at a temperature of 25±3°C for 7 weeks. Weekly measurements of fresh and dry weights of the adventitious roots were taken for analysis of growth rates, and growth curves were constructed using data from two replications. At the 4-week mark of the culture period, 100 μM MeJa, sterilized

through a 0.2- $\mu$ m filter, was introduced into the BTBB culture and incubated for durations of 1, 2, 3, and 4 weeks.

#### Biomass measurement of adventitious roots

The fresh weight of the harvested adventitious roots (1, 2, 3, and 4 weeks of culture) was obtained by gently blotting dry the adventitious roots on a paper towel for 15 min and weighing using an analytical balance. The adventitious roots were dried at room temperature (31-34°C) for 2-3 days to get the dry weight then ground into powder using a pestle and mortar. The growth index of adventitious root cultures was calculated using equation (1):

$$\text{Growth index} = \frac{\text{Final dry weight} - \text{Initial dry weight}}{\text{Initial dry weight}} \quad (1)$$

The growth rate of the cultured adventitious roots was calculated using equation (2):

$$\text{Growth rate} = \frac{\ln(\text{final dry weight}) - \ln(\text{initial dry weight})}{\text{length of culture periods (days)}} \quad (2)$$

#### Quantification of total phenolic and flavonoid contents

The maceration method was used to extract adventitious roots to get the total phenolic compound and flavonoid. One gram (1g) of adventitious root powder was mixed with 20 mL methanol and then incubated for 48 hours in a shaker incubator at 7-10 rpm. Using filter paper the liquid extract was separated from solid debris and concentrated by evaporation to 2 mL at room temperature. For phenolic compound measurement, 200  $\mu$ L of the methanol extract was mixed with 1 mL of 1:10 Folin-Ciocalteu reagents and incubated for 5 minutes at room temperature. Next, the solution was added 0.8 mL of 7.5% (w/v) sodium carbonate ( $\text{Na}_2\text{CO}_3$ ) then incubated for 30 minutes at room temperature. The absorbance of the solution was measured at 765 nm using a UV spectrophotometer (BOECO S-22, Germany). Determining the total phenolic compounds was done by regression analysis from the absorbance rate of standard phenolic acid. Results were expressed as mg GEA (gallic acid equivalent) per gram of dry weight (Kaewseejan et al., 2015).

The concentration of flavonoids was measured using 250  $\mu\text{L}$  of the methanol extract that was mixed with 125 mL distilled water and 75  $\mu\text{L}$  of 5% (w/v) sodium nitrate, then the mixture was incubated at room temperature for 6 minutes. Next, the solution was added to 150  $\mu\text{L}$  of 10% (w/v) aluminum chloride and then incubated for 5 minutes. One hundred and fifty microlitres of sodium hydroxide were added and topped up with distilled water to 2.5 mL total volume. The absorbance of the solution was measured using a UV spectrophotometer at 510 nm. Determining the flavonoid concentration was done by regression analysis from the absorbance rate of standard quercetin and kaempferol (Kaewseejan et al., 2015).

#### Malondialdehyde and proline measurement

MDA and proline levels in MeJa-treated adventitious root cultures were determined following the procedure outlined by Cui et al., 2010. Cui et al., 2010. Initially, 0.2 g of fresh adventitious roots were homogenized by adding 4 mL of 0.1% (w/v) trichloroacetic acid (TCA) using a mortar and pestle. The resulting mixture was then subjected to centrifugation at 3,000 rpm for 10 minutes. The supernatant obtained was combined with 0.25 mL of 200% TCA, 0.25 mL of butylated hydroxytoluene (BHT), and 0.5 mL of 0.65% (w/v) thiobarbituric acid (TBA). This mixture was incubated at 95°C in a water bath for 30 minutes, followed by rapid cooling on ice. Subsequently, the solution was centrifuged again at 3,000 rpm for 10 minutes. The absorbance of the resulting supernatant was measured using a UV spectrophotometer (Multiscan Go - Thermosci) at wavelengths of 523 nm and 600 nm. The concentration of MDA was calculated using equation (3).

$$\text{Malondialdehyde (MDA)} = \frac{\text{Absorbance value at } \lambda_{523} - \text{Absorbance value at } \lambda_{600}}{155 \text{ mM}^{-1}\text{cm}^{-1}(\text{Coefficient value MDA-TBA})} \quad (3)$$

Lipid peroxidation was calculated according to Puangpronpitag et al. (2010). Puangpronpitag et al., (2010). The absorbance of MDA for each sample ( $A_{\text{sample}}$ ) and phosphate buffer ( $A_{\text{control}}$ ) was measured at 532 nm and calculated using equation (4):

$$\text{Inhibition (\%)} = \frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \cdot 100\% \quad (4)$$

The concentration of proline was performed according to Bates et al., 1973. First, 0.5 g of fresh adventitious roots were homogenized using a mortar and pestle while adding 10 ml of 3% (v/v) sulfosalicylic acid. The mixture was centrifuged at 6,000 rpm for 10 min. The supernatant was mixed with 2 mL ninhydrin acid and 2 ml glacial acetate before being incubated at 100°C in a water bath for 1 hour, followed by immediate cooling on ice. Next, the solution was added 4 mL of toluene and mixed thoroughly by vortexing for 15-20 seconds. The chromophore was measured using a UV spectrophotometer at 520 nm with pure toluene as a blank solution. Proline concentration was calculated based on the L-proline standard solution.

Measurement of physical and chemical conditions of the media

The changes of pH, total sugar level (Atago, Master 10T), and electrical conductivity (Ezdo, Condo5021) of culture media were determined at different stages of the experiment: pre- and post-sterilization and at the end of the culture period.

## RESULT


Quantitative analysis of *G. procumbens* adventitious roots growth rate

Adventitious roots derived from both leaves and stems of *G. procumbens* were cultured and maintained in a BTBB. The weights of the adventitious roots, both fresh and dry, were documented at 7-day intervals to ascertain the appropriate starting time for MeJa elicitation (see Figure 1). During the initial two weeks of culture, the growth of adventitious root cultures was sluggish, indicating a lag phase where the initial explants were acclimating to the culture conditions. Subsequently, a notable surge in both fresh and dry weights was observed in the third and fourth weeks of culture, signifying a logarithmic or exponential growth phase.

However, by the fifth week of culture, the rate of growth of the adventitious roots decelerated, indicating a transition towards a stationary phase of growth. The peak biomass was recorded in the sixth week of culture, yielding 35.8 g of fresh weight and 2.005 g of dry




weight. After the sixth week, both fresh and dry weights of adventitious roots exhibited a decline. This growth pattern aligns with previous reports on the hairy root culture of *Glycyrrhiza* reported by Wongwicha et al. (2011) and Wongwicha et al., (2011). Given that MeJa elicitation is recommended during the exponential growth phase, we introduced the MeJa elicitor in the fourth week of culture for subsequent experiments.




**Figure 1.** The growth trajectory of *G. procumbens* adventitious root culture within BTBBs.

The effect of different MeJa elicitation periods on the growth of *G. procumbens* adventitious roots

After adding MeJa to the culture medium, the fresh and dry weights of *G. procumbens* adventitious roots were increased from 0 days (M0) to the third week of elicitation (M3) (Figure 2A). These results showed that MeJa has a positive effect on the biomass of *G. procumbens* adventitious roots. The growth index of MeJa-treated *G. procumbens* adventitious roots, however, was negatively correlated with its growth rate (Figure 2B).



(A)




(B)

**Figure 2.** *G. procumbens* adventitious root growth elicited with 100  $\mu$ M MeJa at different periods (A: Fresh and dry weights of MeJa-treated adventitious roots at different elicitation periods; B: Growth index and growth rate of MeJa-treated adventitious roots at different elicitation periods; M0: 0 days of MeJa elicitation (4 week-old culture); M1-M4: one to four weeks of MeJa elicitation)

The effect of the MeJa elicitation period on phenol and flavonoid production

To assess the impact of MeJA on the production of phenolic and flavonoid compounds, a concentration of 100  $\mu$ M MeJa was administered to 4-week-old adventitious root cultures. Our findings revealed that MeJa elicitation significantly heightened the accumulation of


phenolic and flavonoid compounds, including <sup>5</sup> kaempferol and quercetin, within *G. procumbens* adventitious roots in the BTBB (see Figure 3). The peak concentrations of phenolic compound (484.49 mg/L/10 g DW), kaempferol (3137.29 mg/L/10 g DW), and quercetin (882.67 mg/L/10 g DW) were noted during the third week of MeJa elicitation (see Figure 3).




**Figure 3.** The concentration of phenolic and flavonoids (kaempferol and quercetin) in MeJa-treated (Each data point represents the mean  $\pm$  standard error value (n=2))

The effect of MeJa elicitation periods on the concentration of <sup>4</sup> MDA and proline

The concentration of <sup>3</sup> MDA and proline of MeJa-treated adventitious root cultures was measured to examine the level of cellular stress. We found that the highest level of MDA (80432.61 nmol/0.5 g FW) and proline (6665.32 nmol/0.5 g FW) was recorded after 1 week of elicitation (Figure 4). The concentration of MDA and proline in MeJa-treated adventitious root cultures after 1, 2, 3, and 4 weeks of elicitation was generally higher than 0 days.




(A)



(B)

**Figure 4.** The concentration of MDA and proline in *G. procumbens* adventitious root cultures treated with MeJa for different elicitation periods

In this study, during elicitation from the start culture, until week four (M0 – M4) MDA and proline profiles have the same patterns, except in week three (M3). This is due to the lack of repetitions so less data is obtained. Accumulation of MDA in the adventitious root of *G. procumbens* was followed by the accumulation of proline. This result indicated that the accumulation of proline was a response of MeJa treatments to peroxidation lipid activity. This accumulation was triggered by a higher concentration of MeJa because proline accumulation is found when plants get various kinds of abiotic stress (Anwar Hossain et al., 2014). Most previous studies have reported that the concentration of proline usually increases when plants have abiotic stresses (Yue et al., 2019).



**Figure 5.** The overview of culture protocol for *G. procumbens* adventitious roots in BTBB (A: Leaf and stem explants of *G. procumbens* were used to induce adventitious roots; B: Adventitious root cultures were cultured in liquid MS medium supplemented with 5 mg/L IBA; C: Adventitious root cultures were transferred to BTBB and incubated in liquid MS medium supplemented with 100  $\mu$ M MeJa (right) and without MeJa (left); D: Fresh (left) and dry (right) weights of adventitious roots collected from BTBB)

This study has established a culture protocol for *G. procumbens* adventitious roots in BTBB (Figure 5). The culture was started to induce adventitious roots from leaves explant in MS solid medium supplemented with 5 mg/L indole-3-butyric acids (IBA), whereas the stem explants were cultured on MS solid medium supplemented with 2 mg/L indole acetic acid (IAA), 4 mg/L 6-benzylaminopurine (BAP), then adventitious roots were culture in Erlenmeyer with 100 mL liquid MS medium supplemented with 5 mg/L IBA. Cultured was transferred to BTBB and the medium was added by various concentrations of MeJa.

## DISCUSSION

Jasmonic acid is an important plant signaling molecule for plant growth and development as well as defense responses. In this study, we added 100  $\mu$ M MeJa based on the previously reported concentration (Jiao et al., 2018; Mangas et al., 2006; Yi et al., 2019) into the culture media containing 4-week-old adventitious roots. We found that the fresh and dry weights of MeJa-treated adventitious root cultures were increased for the first three weeks of culture. Supplementation of MeJa in culture medium has been reported to improve the plant growth in several plant species, such as cell suspension culture of *Melastoma malabathricum* (See et al., 2011), callus culture of Carnation (Matter et al., 2017) and adventitious root culture of *Centella asiatica* (L.) Urb (Nguyen et al., 2019). Nevertheless, MeJa could negatively affect plant growth, as reported in the cell suspension culture of *Thevetia peruviana* (Pers.) K. Schum (Mendoza et al., 2018a), *Taxus* (Patil et al., 2014), and strawberries (Miyanaga et al., 2000).

MeJA has been extensively employed as an elicitor in numerous plant species to augment flavonoid production in cell culture systems. For instance, MeJa has been introduced into culture media to elevate flavonoid production in *Hypericum perforatum* cell cultures (Cui et al., 2010) (Cui et al., 2010), ginsenoside in American ginseng root hair cultures (Kochan et al., 2018) (Kochan et al., 2018), and triterpenoid in *Centella asiatica* (L.) Urb root hair cultures (Nguyen et al., 2019) (Nguyen et al., 2019). In this investigation, we found that the addition of MeJa to the culture media for one to three weeks notably enhanced the biomass of phenolic compounds and flavonoid accumulations in adventitious root cultures of *G. procumbens*. However, a more prolonged exposure time (4 weeks of elicitation) resulted in reduced biomass accumulation and concentration of phenolic compounds and flavonoids. Comparable results were documented in *Thevetia peruviana* cell suspension cultures, where the highest phenol and flavonoid concentrations were achieved after 72 and 96 hours of MeJa elicitation (Mendoza et al., 2018) (Mendoza et al., 2018). Extended exposure times (7 days) with a high concentration of MeJa were observed to enhance the production of saponin and ginsenoside in American ginseng root hair cultures (Kochan et al., 2018), while a lower MeJa concentration (100  $\mu$ M) increased the accumulation of ginsenoside in *Panax ginseng* adventitious roots (Kim et al., 2004). Prolonged exposure to MeJa has been reported to inhibit the production of secondary

metabolites in *Centella asiatica* (L.) Urb. root hairy cultures (Nguyen et al., 2019) (Nguyen et al., 2019) and *Thevetia peruviana* cell suspension cultures (Mendoza et al., 2018) (Mendoza et al., 2018 ). This adverse effect may be attributed to the direct toxic impact of MeJA or a loss of culture viability.

In our research, we observed an increase in both MDA and proline levels following MeJA elicitation. Adventitious root cultures can face stress due to the high levels of agitation and aeration present in an *in vitro* system like the BTBB. This *in vitro* environment induces oxidative stress in plant cells due to the heightened presence of reactive oxygen species (ROS). ROS compounds are detrimental to plant cells and can lead to the degradation of lipid membranes, proteins, and DNA (Davey et al., 2005; Esfandiari et al., 2007). (Davey et al., 2005; Esfandiari et al., 2007). MDA, a byproduct of lipid peroxidation caused by ROS, serves as a marker for oxidative stress. An elevation in MDA concentration was noted in adventitious roots of *Hypericum perforatum* when cultured in MS media containing a high nitrate to ammonium ratio and high sucrose concentration (5-9%) (Baque et al., 2014). (Baque et al., 2014)

Proline, a proteinogenic amino acid crucial for plant primary metabolism, acts as an osmolyte protectant. Additionally, proline stabilizes proteins and mitigates the detrimental effects of ROS under stressful conditions (Kishor, 2015) (Kishor, 2015). It has been documented that proline accumulates in liquid culture media when plant cells or tissues are exposed to mineral stress (Baque et al., 2014) (Baque et al., 2014), osmotic stress (Lee & Paek, 2012) (Lee & Paek, 2012), or addition of hormones into the media. Furthermore, a high concentration of inorganic salts in culture media resulted in increased proline concentration in *Spathiphyllum cannifolium* (Dewir et al., 2005) (Dewir et al., 2005).

## CONCLUSION

Overall, we have established a simple and robust *in vitro* system for *G. procumbens* adventitious roots using BTBB. The growth of adventitious roots of *G. procumbens* and its biomass were influenced by MeJA elicitation. The highest biomass was attained at 3 weeks

of MeJa elicitation compared to six weeks of culture without MeJa treatment in BTBB. The concentration of phenol, kaempferol, and quercetin MeJa-treated adventitious root cultures were 2.2-, 2.6-, and 3.5-fold higher than non-treated control after 3 weeks of culture. MDA and proline concentrations in MeJa-treated adventitious root cultures were higher than in non-treated control, indicating the importance of MeJa in alleviating oxidative stress. The pH, total sugar level, and electrical conductivity of the culture media were decreased at the end of the culture period. Our research outcomes carry significant implications for forthcoming biotechnological applications, particularly for large-scale production of valuable bioactive compounds intended for industrial use.

#### **ACKNOWLEDGMENTS**

This research was funded by the Directorate Research and Community Service, Ministry of Research, Technology and Higher Education, Indonesia, Dissertation Research with grand number 8/E1/KPT/2021; contract number 4/E1/KP.PTNBH/2021 and 499/UN3.15/PT/2021.

#### **REFERENCES**

- Algariri, K., Meng, K. Y., Atangwho, I. J., Asmawi, M. Z., Sadikun, A., Murugaiyah, V., & Ismail, N. (2013). Hypoglycemic and anti-hyperglycemic study of *Gynura procumbens* leaf extracts. *Asian Pacific Journal of Tropical Biomedicine*, 3(5), 358–366. [https://doi.org/10.1016/S2221-1691\(13\)60077-5](https://doi.org/10.1016/S2221-1691(13)60077-5)
- Ali, M., Hahn, E.-J., & Paek, K.-Y. (2007). Methyl Jasmonate and Salicylic Acid Induced Oxidative Stress and Accumulation of Phenolics in *Panax ginseng* Bioreactor Root Suspension Cultures. *Molecules*, 12(3), 607–621. <https://doi.org/10.3390/12030607>
- Anwar Hossain, M., Hoque, M. A., Burritt, D. J., & Fujita, M. (2014). Proline Protects Plants Against Abiotic Oxidative Stress: Biochemical and Molecular Mechanisms. In *Oxidative Damage to Plants: Antioxidant Networks and Signaling* (pp. 477–522). Elsevier Inc. <https://doi.org/10.1016/B978-0-12-799963-0.00016-2>
- Baque, A., Murthy, H. N., & Paek, K.-Y. (2014). Adventitious Root Culture of *Morinda citrifolia* in Bioreactors for Production of Bioactive Compounds. In *Production of Biomass and Bioactive Compounds Using Bioreactor Technology* (pp. 185–222). Springer Netherlands. [https://doi.org/10.1007/978-94-017-9223-3\\_9](https://doi.org/10.1007/978-94-017-9223-3_9)



- Baque, Md. A., Shiragi, Md. H. K., Moh, S.-H., Lee, E.-J., & Paek, K.-Y. (2013). Production of biomass and bioactive compounds by adventitious root suspension cultures of *Morinda citrifolia* (L.) in a liquid-phase airlift balloon-type bioreactor. *In Vitro Cellular & Developmental Biology - Plant*, 49(6), 737–749. <https://doi.org/10.1007/s11627-013-9555-3>
- Bates, L. S., Waldren, R. P., & Teare, I. D. (1973). Rapid determination of free proline for water-stress studies. *Plant and Soil*, 39(1), 205–207. <https://doi.org/10.1007/BF00018060>
- Choi, S. M., Ho Son, S., Rho Yun, S., Woung Kwon, O., Hoon Seon, J., & Yoeup Paek, K. (2000). Pilot-Scale Culture of Adventitious Roots of Ginseng in a Bioreactor System. *Plant Cell, Tissue and Organ Culture*, 62(3), 187–193. <https://doi.org/10.1023/A:1006412203197>
- Cui, X.-H., Chakrabarty, D., Lee, E.-J., & Paek, K.-Y. (2010). Production of adventitious roots and secondary metabolites by *Hypericum perforatum* L. in a bioreactor. *Bioresource Technology*, 101(12), 4708–4716. <https://doi.org/10.1016/j.biortech.2010.01.115>
- Davey, M. W., Stals, E., Panis, B., Keulemans, J., & Swennen, R. L. (2005). High-throughput determination of malondialdehyde in plant tissues. *Analytical Biochemistry*, 347(2), 201–207. <https://doi.org/10.1016/j.ab.2005.09.041>
- Dewir, Y. H., Chakrabarty, D., Ali, M. B., Hahn, E. J., & Paek, K. Y. (2005). Effects of Hydroponic Solution EC, Substrates, PPF and Nutrient Scheduling on Growth and Photosynthetic Competence During Acclimatization of Micropropagated *Spathiphyllum* plantlets. *Plant Growth Regulation*, 46(3), 241–251. <https://doi.org/10.1007/s10725-005-0161-1>
- Esfandiari, E., Shekari, F., Shekari, F., & Esfandiari, M. (2007). The effect of salt stress on antioxidant enzymes activity and lipid peroxidation on the wheat seedling. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 35(1), 48–56.
- Faizah, H., Tanjung, M., Purnobasuki, H., & Manuhara, Y. S. W. (2018). Biomass and flavonoid production of *Gynura procumbens* (L.) merr adventitious root culture in balloon-type bubble-bioreactor influenced by elicitation. *Asian Journal of Plant Sciences*, 17(2), 107–119. <https://doi.org/10.3923/AJPS.2018.107.119>
- Foyer, C. H., & Harbinson, J. (2019). Oxygen Metabolism and the Regulation of Photosynthetic Electron Transport. In *Causes of Photooxidative Stress and Amelioration of Defense Systems in Plants* (pp. 1–42). CRC Press. <https://doi.org/10.1201/9781351070454-1>
- Ho, T.-T., Murthy, H. N., & Park, S.-Y. (2020). Methyl Jasmonate Induced Oxidative Stress and Accumulation of Secondary Metabolites in Plant Cell and Organ Cultures.

- International Journal of Molecular Sciences*, 21(3), 716.  
<https://doi.org/10.3390/ijms21030716>
- Hoe, S.-Z., Lee, C.-N., Mok, S.-L., Kamaruddin, M. Y., & Lam, S.-K. (2011). *Gynura procumbens* Merr. decreases blood pressure in rats by vasodilatation via inhibition of calcium channels. *Clinics*, 66(1), 143–150. <https://doi.org/10.1590/S1807-59322011000100025>
- Iskander, M. N., Song, Y., Coupar, I. M., & Jiratchariyakul, W. (2002). Antiinflammatory screening of the medicinal plant *Gynura procumbens*. *Plant Foods for Human Nutrition*, 57(3/4), 233–244. <https://doi.org/10.1023/A:1021851230890>
- Jarikasem, S., Charuwichitratana, S., Siritantikorn, S., Chantratita, W., Iskander, M., Frahm, A. W., & Jiratchariyakul, W. (2013). Antiherpetic Effects of *Gynura procumbens*. *Evidence-Based Complementary and Alternative Medicine*, 2013, 1–10. <https://doi.org/10.1155/2013/394865>
- Jiang, X.-L., Piao, X.-C., Ri Gao, Jin, M.-Y., Jiang, J., Jin, X.-H., & Lian, M.-L. (2017). Improvement of bioactive compound accumulation in adventitious root cultures of an endangered plant species, *Oplopanax elatus*. *Acta Physiologiae Plantarum*, 39(10), 226. <https://doi.org/10.1007/s11738-017-2525-3>
- Jiao, C., Song, C., Zheng, S., Zhu, Y., Jin, Q., Cai, Y., & Lin, Y. (2018). Metabolic Profiling of *Dendrobium officinale* in Response to Precursors and Methyl Jasmonate. *International Journal of Molecular Sciences*, 19(3), 728. <https://doi.org/10.3390/ijms19030728>
- Kaewseejan, N., Sutthikhum, V., & Siriamornpun, S. (2015). Potential of *Gynura procumbens* leaves as source of flavonoid-enriched fractions with enhanced antioxidant capacity. *Journal of Functional Foods*, 12, 120–128. <https://doi.org/10.1016/j.jff.2014.11.001>
- Karuppusamy, S. (2009). A review on trends in production of secondary metabolites from higher plants by in vitro tissue, organ and cell cultures. In *Journal of Medicinal Plants Research* (13th ed., Vol. 3, pp. 1222–1239).
- Kavi Kishor, P. B. (2015). Role of proline in cell wall synthesis and plant development and its implications in plant ontogeny. *Frontiers in Plant Science*, 6. <https://doi.org/10.3389/fpls.2015.00544>
- Kim, J., Lee, C.-W., Kim, E. K., Lee, S.-J., Park, N.-H., Kim, H.-S., Kim, H.-K., Char, K., Jang, Y. P., & Kim, J.-W. (2011). Inhibition effect of *Gynura procumbens* extract on UV-B-induced matrix-metalloproteinase expression in human dermal fibroblasts. *Journal of Ethnopharmacology*, 137(1), 427–433. <https://doi.org/10.1016/j.jep.2011.04.072>
- Kim, Y., Hahn, E., Murthy, H. N., & Paek, K. (2004). Adventitious root growth and ginsenoside accumulation in *Panax ginseng* cultures as affected by methyl jasmonate.

- Biotechnology Letters*, 26(21), 1619–1622. <https://doi.org/10.1007/s10529-004-3183-2>
- Kochan, E., Szymczyk, P., Kuźma, Ł., Szymańska, G., Wajs-Bonikowska, A., Bonikowski, R., & Sienkiewicz, M. (2018). The Increase of Triterpene Saponin Production Induced by Trans-Anethole in Hairy Root Cultures of *Panax quinquefolium*. *Molecules*, 23(10), 2674. <https://doi.org/10.3390/molecules23102674>
- Krishnan, V., Ahmad, S., & Mahmood, M. (2015). Antioxidant Potential in Different Parts and Callus of *Gynura procumbens* and Different Parts of *Gynura bicolor*. *BioMed Research International*, 2015, 1–7. <https://doi.org/10.1155/2015/147909>
- Lee, E. J., Park, S. Y., & Paek, K. Y. (2015). Enhancement strategies of bioactive compound production in adventitious root cultures of *Eleutherococcus koreanum* Nakai subjected to methyl jasmonate and salicylic acid elicitation through airlift bioreactors. *Plant Cell, Tissue and Organ Culture (PCTOC)*, 120(1), 1–10. <https://doi.org/10.1007/s11240-014-0567-4>
- Lee, E.-J., Moh, S.-H., & Park, S.-Y. (2014). Production of Biomass and Bioactive Compounds in Adventitious Root Cultures of *Eleutherococcus koreanum* Nakai. In *Production of Biomass and Bioactive Compounds Using Bioreactor Technology* (pp. 223–249). Springer Netherlands. [https://doi.org/10.1007/978-94-017-9223-3\\_10](https://doi.org/10.1007/978-94-017-9223-3_10)
- Lee, E.-J., & Paek, K.-Y. (2012). Effect of nitrogen source on biomass and bioactive compound production in submerged cultures of *Eleutherococcus koreanum* nakai adventitious roots. *Biotechnology Progress*, 28(2), 508–514. <https://doi.org/10.1002/btpr.1506>
- Mangas, S., Bonfill, M., Osuna, L., Moyano, E., Tortoriello, J., Cusido, R. M., Teresa Piñol, M., & Palazón, J. (2006). The effect of methyl jasmonate on triterpene and sterol metabolisms of *Centella asiatica*, *Ruscus aculeatus* and *Galphimia glauca* cultured plants. *Phytochemistry*, 67(18), 2041–2049. <https://doi.org/10.1016/j.phytochem.2006.06.025>
- Manogaran, M., Lim, V., & Mohamed, R. (2019). Phytoconstituents of the *Gynura procumbens* ethanol leaf extract and its fractions and their effects on viability of macrophages. *Journal of Herbmed Pharmacology*, 8(3), 224–230. <https://doi.org/10.15171/jhp.2019.33>
- Manuhara, Y. S. W., Kusuma, D. Y., Sari, R. L. K., & Kristanti, A. N. (2017). Biomass Production of *Gynura procumbens* Adventitious Roots in Different Type of Liquid Culture. *Biosaintifika: Journal of Biology & Biology Education*, 9(3), 523–529. <https://doi.org/10.15294/biosaintifika.v9i3.9670>
- Matter, M. A., Hanafy, M. S., & Aly, U. I. (2017). Effect of Methyl Jasmonate and Mannitol Application on Growth and Eugenol Content in Callus Cultures of Carnation. *Plant*

*Tissue Culture and Biotechnology*, 27(2), 227–240.  
<https://doi.org/10.3329/ptcb.v27i2.35028>

- Mendoza, D., Cuaspid, O., Arias, J. P., Ruiz, O., & Arias, M. (2018). Effect of salicylic acid and methyl jasmonate in the production of phenolic compounds in plant cell suspension cultures of *Thevetia peruviana*. *Biotechnology Reports*, 19, e00273. <https://doi.org/10.1016/j.btre.2018.e00273>
- Miyanaaga, K., Seki, M., & Furusaki, S. (2000). Quantitative determination of cultured strawberry-cell heterogeneity by image analysis: effects of medium modification on anthocyanin accumulation. *Biochemical Engineering Journal*, 5(3), 201–207. [https://doi.org/10.1016/S1369-703X\(00\)00059-0](https://doi.org/10.1016/S1369-703X(00)00059-0)
- Murthy, H. N., Dandin, V. S., & Paek, K.-Y. (2016). Tools for biotechnological production of useful phytochemicals from adventitious root cultures. *Phytochemistry Reviews*, 15(1), 129–145. <https://doi.org/10.1007/s11101-014-9391-z>
- Muthoharoh, L., Faizah, H., Hartatie Hardjo, P., Novi Kristanti, A., & Sri Wulan Manuhara, Y. (2019). Effect of Carbon Source on Biomass and Flavonoid Content of *Gynuraprocumbens* (Lour.) Merr Adventitious Root in Liquid Culture. *Biosciences, Biotechnology Research Asia*, 16(1), 121–127. <https://doi.org/10.13005/bbra/2729>
- Ng, H.-K., Poh, T.-F., Lam, S.-K., & Hoe, S.-Z. (2013). Potassium channel openers and prostacyclin play a crucial role in mediating the vasorelaxant activity of *Gynura procumbens*. *BMC Complementary and Alternative Medicine*, 13(1), 188. <https://doi.org/10.1186/1472-6882-13-188>
- Nguyen, K. Van, Pongkitwitoon, B., Pathomwichaiwat, T., Viboonjun, U., & Prathantururug, S. (2019). Effects of methyl jasmonate on the growth and triterpenoid production of diploid and tetraploid *Centella asiatica* (L.) Urb. hairy root cultures. *Scientific Reports*, 9(1), 18665. <https://doi.org/10.1038/s41598-019-54460-z>
- Nurulita, N. A., Meiyanto, E., Sugiyanto, Matsuda, E., & Kawaichi, M. (2012). *Gynura procumbens* modulates the microtubules integrity and enhances distinct mechanism on doxorubicin and 5-fluorouracil-induced breast cancer cell death. *Oriental Pharmacy and Experimental Medicine*, 12(3), 205–218. <https://doi.org/10.1007/s13596-012-0063-5>
- Paek, K.-Y., Hahn, E.-J., & Son, S.-H. (2001). Application of bioreactors for large-scale micropropagation systems of plants. *In Vitro Cellular & Developmental Biology - Plant*, 37(2), 149–157. <https://doi.org/10.1007/s11627-001-0027-9>
- Patil, R. A., Lenka, S. K., Normanly, J., Walker, E. L., & Roberts, S. C. (2014). Methyl jasmonate represses growth and affects cell cycle progression in cultured *Taxus* cells. *Plant Cell Reports*, 33(9), 1479–1492. <https://doi.org/10.1007/s00299-014-1632-5>
- Puangpronpitag, D., Chaichanadee, S., Naowaratwattana, W., & Al, E. (2010). *Gynura* .. extract.pdf. In *Asian Journal of Plant Sciences* 9 (3): 146-151, 2010 ISSN 1682-3974 (Vol. 9, Issue 3).

- Rahimi, S., Kim, Y.-J., & Yang, D.-C. (2015). Production of ginseng saponins: elicitation strategy and signal transductions. *Applied Microbiology and Biotechnology*, *99*(17), 6987–6996. <https://doi.org/10.1007/s00253-015-6806-8>
- Saiman, M. Z., Mustafa, N. R., Schulte, A. E., Verpoorte, R., & Choi, Y. H. (2012). Induction, characterization, and NMR-based metabolic profiling of adventitious root cultures from leaf explants of *Gynura procumbens*. *Plant Cell, Tissue and Organ Culture (PCTOC)*, *109*(3), 465–475. <https://doi.org/10.1007/s11240-011-0111-8>
- Takanashi, K., Dan, K., Matsuoka, T., & Torii, K. (2019). The Cancer Cell Killing Effects of *Gynura procumbens*, Toleaf Are Associated with CD4+ T Lymphocyte Activation. *Pharmaceutical Frontiers*, *:e190004*, 1–13. <https://doi.org/10.20900/pf20190004>
- Tan, H.-L., Chan, K.-G., Pusparajah, P., Lee, L.-H., & Goh, B.-H. (2016). *Gynura procumbens*: An Overview of the Biological Activities. *Frontiers in Pharmacology*, *7*(52). <https://doi.org/10.3389/fphar.2016.00052>
- Wang, Q., Wang, J., Chai, H., Li, J., Man, S., & Gao, W. (2015). Optimization of balloon-type bubble bioreactor angle and methyl jasmonate concentration to enhance metabolite production in adventitious roots of *Pseudostellaria heterophylla*. *Research on Chemical Intermediates*, *41*(8), 5555–5563. <https://doi.org/10.1007/s11164-014-1681-y>
- Wongwicha, W., Tanaka, H., Shoyama, Y., & Putalun, W. (2011). Methyl Jasmonate Elicitation Enhances Glycyrrhizin Production in *Glycyrrhiza inflata* Hairy Roots Cultures. *Zeitschrift Für Naturforschung C*, *66*(7–8), 423–428. <https://doi.org/10.1515/znc-2011-7-815>
- Wu, S. Q., Lian, M. L., Gao, R., Park, S. Y., & Piao, X. C. (2011). Bioreactor application on adventitious root culture of *Astragalus membranaceus*. *In Vitro Cellular & Developmental Biology - Plant*, *47*(6), 719–724. <https://doi.org/10.1007/s11627-011-9376-1>
- Yi, T. G., Park, Y., Park, J.-E., & Park, N. II. (2019). Enhancement of Phenolic Compounds and Antioxidative Activities by the Combination of Culture Medium and Methyl Jasmonate Elicitation in Hairy Root Cultures of *Lactuca indica* L. *Natural Product Communications*, *14*(7), 1–9. <https://doi.org/10.1177/1934578X19861867>
- Yue, Y., Jiang, H., Du, J., Shi, L., Bin, Q., Yang, X., & Wang, L. (2019). Variations in physiological response and expression profiles of proline metabolism-related genes and heat shock transcription factor genes in petunia subjected to heat stress. *Scientia Horticulturae*, *258*(108787), 1–10. <https://doi.org/10.1016/j.scienta.2019.108811>

## ORIGINALITY REPORT

11%

SIMILARITY INDEX

13%

INTERNET SOURCES

15%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://link.springer.com">link.springer.com</a> Internet Source	2%
2	<a href="http://mail.scialert.net">mail.scialert.net</a> Internet Source	2%
3	<a href="http://journal.ipb.ac.id">journal.ipb.ac.id</a> Internet Source	2%
4	<a href="http://journal.ugm.ac.id">journal.ugm.ac.id</a> Internet Source	1%
5	<a href="http://repository.ubaya.ac.id">repository.ubaya.ac.id</a> Internet Source	1%
6	<a href="http://repository.uam.ac.id">repository.uam.ac.id</a> Internet Source	1%
7	Xi-Hua Cui, Hosakatte Niranjana Murthy, Chun-Hua Wu, Kee-Yoeup Paek. "Sucrose-induced osmotic stress affects biomass, metabolite, and antioxidant levels in root suspension cultures of <i>Hypericum perforatum</i> L.", <i>Plant Cell, Tissue and Organ Culture (PCTOC)</i> , 2010 Publication	1%

8

"Date Palm Biotechnology Protocols Volume I", Springer Science and Business Media LLC, 2017

Publication

1 %

9

"Nutraceuticals Production from Plant Cell Factory", Springer Science and Business Media LLC, 2022

Publication

1 %

Exclude quotes Off

Exclude matches < 1%

Exclude bibliography On