

Fermentation quality of king grass (*Pennisetum purpureophoides*) ensiled with epiphytic lactic acid bacteria and tannin of acacia¹

B. Santoso,* B. Tj. Hariadi,* H. Manik† and H. Abubakar‡

*Department of animal Nutrition, Faculty of Animal Science, Fishery and Marine Science, State University of Papua, Indonesia; †Department of Animal Production, Faculty of Animal science, Fishery and Marine science, State University of Papua, Indonesia; ‡Department of Biology, Faculty of Mathematics and Natural Science, State University of Papua

ABSTRACT: The aim of this study was to evaluate the fermentation quality of king grass (*Pennisetum purpureophoides*) ensiled with additions of epiphytic lactic acid bacteria (LAB) prepared from fermented grass extract (FGE) or combined with tannin of acacia. Six treatments were (A) king grass without additive as a control; (B) king grass + 3% (v/w) of FGE; (C) king grass + 3% (v/w) of FGE + 10 ml of acacia extract (50 g/100 ml); (D) king grass + 3% of FGE (v/w) + 10 ml of acacia extract (50 g/75 ml); (E) king grass + 3% of FGE (v/w) + 10 ml of acacia extract (50 g/50 ml), and (F) king grass + 3% of FGE (v/w) + 10 ml of acacia extract (50 g/25 ml). About 250 g of silage materials were ensiled in 400 ml bottle silos at room temperatures (approximately 28 °C) for 30 days. Results showed that the number of lactic acid bacteria in FGE was increased from 77×10^5 cfu/ml to 286×10^5 cfu/ml after 2 days anaerobic incubation. Concentration of lactic acid in silages with addition of FGE or combined with tannin of acacia (B, C, D, E and F) were higher ($P < 0.01$) than that of silage A (control). Silages with addition of FGE combined with tannin of acacia (C, D, E and F) had lower pH value than that of silages A and B. Concentrations of $\text{NH}_3\text{-N}$ was decreased with increasing concentration of tannin. Butyric acid concentration was decreased in silages of B, C, D, E and F as compared to silage A. It was concluded that addition of FGE combined with tannin prepared from acacia leaf improved fermentation quality of king grass silage.

Key words: silage, grass, lactic acid, tannin

INTRODUCTION

It is recognized that tropical grasses have low water soluble carbohydrate content, high buffering capacity and low lactic acid bacteria (LAB) number (Smith, 1962; McDonald et al., 1991; Yahaya et al., 2004b). These properties result in low lactic acid production; hence it is difficult to produce good-quality silage from tropical grasses. The epiphytic microorganisms existed naturally in forage crops are responsible for silage fermentation and also influence the effectiveness of silage bacterial inoculation. However, the number of LAB is usually low and vary depended on growing crops (Muck, 1990; Lin et al., 1992).

Fermented grass extract of epiphytic lactic acid bacteria (FGE) is prepared by culturing microorganism adherent to the grass materials before preparation of silage and the grown microorganism are used as a starter of silage fermentation. Ohshima et al. (1997) reported that addition of forage extract that incubated anaerobically for 2 days improved fermentation quality of alfalfa silage. Previous studies of Yahaya et al. (2004b); Bureenok et al. (2006); Masuko et al. (2002) found that addition of fermented grass extract on silage materials could increase lactic acid production, decrease NH_3N concentration, increase lactic acid bacteria population, and improve fermentation quality of grass silage. Wang et al. (2009) concluded that fermented grass extract can be used as a good source of LAB for direct cut alfalfa silage.

During the ensiling process of forage, extensive proteolysis occurs due to the combined action of both plant and microbial enzymes resulting in conversion of most protein to non-protein nitrogen (NPN) fractions mainly amino acid N, peptide N, and ammonia N (Ohshima and McDonald, 1978;

¹ The research grant was provided by the State University of Papua (Contract No. 037/SP2H/PP/D2M/III/2009). The authors are grateful to Nurhaida K. Tero for technical assistant during experiment.

Harrison and Blauwiel, 1994). The rapid rate of degradation of silage NPN and soluble-protein N in the rumen results in a pronounced peak in rumen ammonia concentration following ingestion (Thomas and Thomas, 1985). Excess quantities of ammonia in the rumen is absorbed into the blood stream, converted to urea in liver and subsequently excreted in the urine, contributing to environmental pollution (Tamminga, 1992).

Tannins are polyphenolic compounds of plant origin that have ability to bind protein. It has been reported that tannin-containing species such as quebracho, chestnut, residues of green tea and black tea reduced protein degradation during ensiling and protein disappearance in the rumen (Salawu et al., 1999; Tobacco et al., 2006; Kondo et al., 2004; Santoso et al., 2007) by inhibiting plant and microbial enzymes or by forming complexes with protein (McSweeney et al., 2001). Hariadi and Santoso (2010) reported that acacia leaf (*Acacia mangium* Willd) contains 5.4% tannin total, hence it is potentially used as silage additive to protect protein degradation during ensiling.

The objective of this experiment was to evaluate the fermentation quality of king grass (*Pennisetum purpureophoides*) ensiled with addition of epiphytic LAB prepared from fermented grass extract or combined with tannin of acacia.

MATERIALS AND METHODS

Forage Material

King grass (*Pennisetum purpureophoides*) was planted at in a 9 m² plot without fertilizer at the experimental field of Faculty of Animal Science, Fishery and Marine Science, State University of Papua in Manokwari, Indonesia. Grass was harvested with a hand clipper in May 2009 after 50 days of regrowth defoliation. The experimental field is located at 134°04' longitude and 00°48' latitude. The area is located at an altitude of 110 m above sea level. The mean annual rainfall and temperature were 159.9 mm and 27.1° C, respectively.

Preparations of Fermented Grass Extract and Acacia Extract

Preparation of FGE according to modified of Bureenok et al. (2006) procedure as previously described by Santoso et al. (2009). The FGE was prepared using 220 g of fresh king grass, which was macerated in 1000 ml of distilled water using a high-speed blender for 4 min. The macerated was filtered through two layers of cheesecloths, and 600 ml of filtrate was collected in Erlenmeyer glass containing 18 g of glucose. The filtrate was mixed well and incubated anaerobically for 48 h at 30 °C. At the end of 48 h, FGE was used as source of LAB. The number of LAB in the FGE was counted before the experiments by using de Man, Rogosa, and Sharpe which were incubated for 3 days at 35 °C (Bureenok et al., 2006).

About 50 g of finely acacia leaf were weighed into 100 ml beaker glass and added 25 ml of distilled water. The same procedure was also performed for 50, 75, and 100 ml of distilled water. The mixtures were boiled for 10 min on a hotplate and filtered through 2 layers of cheesecloth. The filtrates were collected and stored at 4 °C for further use.

Silage Preparation and Treatments

The fresh king grass was wilted at room temperature (approximately 28 °C) for 24 h and chopped into 3 – 5 cm. The chopped grass was thoroughly mixed and a representative samples obtained. Total of 6 treatments were as follows (A) king grass without additive as a control; (B) king grass + 3% (v/w) of FGE; (C) king grass + 3% (v/w) of FGE + 10 ml of acacia extract (50 g/100 ml); (D) king grass + 3% of FGE (v/w) + 10 ml of acacia extract (50 g/75 ml); (E) king grass + 3% of FGE (v/w) + 10 ml of acacia extract (50 g/50 ml), and (F) king grass + 3% of FGE (v/w) + 10 ml of acacia extract (50 g/25 ml). Both FGE and acacia extract were sprayed onto silage materials using a hand sprayer and subsequently mixed by hand. Based on the concentration of LAB in FGE, the final application was 5.8×10^6 cfu/g of fresh forage. Tannin concentration calculated in silage C, D, E and F were 2.7, 3.6, 5.2 and 10.3 g/kg, respectively. About 250 g of silage materials were packed into 400 ml

laboratory glass bottle silos. Each treatment was prepared in triplicate and the silos were stored in room temperature for 30 days.

Analytical Procedure

Dried samples were used to determine DM, ash and crude protein (CP) according to procedure of AOAC (1990). Procedure of Van Soest et al. (1991) was used to determined concentrations of NDF, acid detergent fiber (ADF) and acid detergent lignin (ADL). NDF was determined without the use of α -amylase and sodium sulfite.

A 20 g of silage was macerated with 70 ml of distilled water and stored at 4 C for 24 h. It was than homogenized for 15 min using a shaker and filtered through a Whatman No. 1542 filter paper. The filtrate was used for determine of pH, VFAs, lactic acid and NH₃-N. The pH value was determined using a pH meter (Hanna Hi 9025). Concentrations of individual VFAs were analyzed using a gas chromatography (Varian CP-9002 GC, Shimadzu, Japan) equipped with flame ionization detector (FID) and stainless steel column (1500 mm × 3 mm i.d). The pressure of nitrogen was 1.25 kg/cm². The temperature of injector oven, column oven and detector were 220, 130 and 220 °C, respectively. Concentrations of lactic acid and NH₃-N were analyzed according to method of Barker and Summerson (1941); Chaney and Marbach (1962), respectively.

Statistical Analysis

Silage fermentation data were subjected to analysis of variance for a completely randomized design using GLM procedure of SAS (SAS Institute Inc., Cary, NC). Duncan's multiple range test was used to separate treatment means, when probability was less than 0.05.

RESULTS AND DISCUSSION

Characteristics of Fermented Grass Extract

Table 1 depicts the result of pH value and LAB number in the fermented grass extract used as additive at ensiling.

Table 1. The pH value and lactic acid bacteria number in grass extract before and after incubation 48 h

	Before incubation	After incubation
pH	6.71	3.51
LAB ($\times 10^5$ cfu/ml)	77	286

The epiphytic LAB number of the material grass was 77×10^5 cfu/ml fresh matter. After 48 h of incubation at 30 °C, the LAB number in the FGE increased to 286×10^5 cfu/ml. Increased LAB number resulted in high concentration of lactic acid, thereby decreasing in pH value from 6.71 to 3.51. In previous study, Santoso et al. (2009) also reported that pH value of fermented grass extract prepared from king grass reduced from 6.41 to 3.45. The patterns of reduced pH value in fermented extracts of grass or legume, and increased LAB number after 48 h of incubation are agree with previous reports of Nishino and Uchida (1999); Bureenok et al. (2006) and Wang et al. (2009). Nishino and Uchida (1999) revealed that several strains of LAB e.g. *Lactobacillus plantarum*, *Lactobacillus viridescens*, *Lactobacillus fermentum*, *Pediococcus acidilactici* were isolated from Lucerne, timothy and cocksfoot fermented extract. However predominant strains were different depended on the material crop used for fermented extract preparation.

Fermentative Quality of The Silages

Fermentation characteristics of king grass silage added with epiphytic lactic acid bacteria and tannin of acacia are shown in Table 2.

Table 2. Fermentation characteristics of king grass silage added with epiphytic lactic acid bacteria and tannin of acacia

	Silages						SE	P
	A	B	C	D	E	F		
LAB, × 10 ⁵ cfu/ml	233	246	140	108.3	128.3	210.3	40.82	0.14
pH	5.28 ^A	4.67 ^{AB}	4.52 ^B	4.31 ^B	4.61 ^B	4.56 ^B	0.14	<0.01
Lactic acid, g/kg DM	0.6 ^B	43.2 ^A	39.7 ^A	39.2 ^A	42.6 ^A	37.4 ^A	6.23	<0.01
NH ₃ -N, g/kg total N	289.7 ^a	159.7 ^b	132.2 ^b	124.0 ^b	120.1 ^b	105.8 ^b	39.30	0.05
Acetic acid, g/kg DM	73.0	95.2	110.0	44.1	75.8	69.2	29.51	0.71
Propionic acid, g/kg DM	11.6	3.9	9.6	6.2	6.3	0	3.50	0.30
Butyric acid, g/kg DM	25.6 ^A	4.7 ^B	4.0 ^B	6.3 ^B	6.7 ^B	5.6 ^B	3.06	0.01
Total VFA, g/kg DM	110.2	103.8	123.6	56.5	88.7	74.8	27.15	0.56

^{a-b} Means in the same row followed by different letters are different (P<0.05)

^{A-B} Means in the same row followed by different letters are different (P<0.01)

The pH values in silages of C, D, E and F were lower (P<0.01) than that of silage A (control). Lower pH value in silage treated with epiphytic LAB and acacia tannin could be due to higher lactic acid concentration in those silages. Seglar (2003) stated that lactic acid is the strongest of all silage acids and its presence will drop pH more effectively than the other volatile fatty acids. Even though, lower pH value found in silage treated with epiphytic LAB, however, the final pH value except silage D are still above than ideal silage pH of 4.0 to 4.5. As reported by Chamberlain and Wilkinson (1996) that secondary fermentation occurs when insufficient acid is produced by primary fermentation to reduce the pH to below a critical level of about 4.5.

Concentration of lactic acid in silages of B, C, D, E and F were significantly (P<0.01) higher as compared to control silage (A). This result agrees with previous study on silages prepared from tropical grasses as reported by Yahaya et al. (2004a); Yahaya et al. (2004b); Bureenok et al. (2006), and Lucerne by Nishino and Uchida (1999); Wang et al. (2009). Increased lactic acid concentration in silage treated with epiphytic LAB could be due to increasing fermentation process by LAB which convert monosaccharide such as glucose and fructose to lactic acid. Higher concentration of lactic acid in silage has advantage to animal. This is because lactic acid can be converted to propionic acid. Chamberlain (1987) reported that lactic acid of silage in the rumen is metabolized via acrylate pathway by *Megasphaera elsdinii*. Furthermore, propionic acid is absorbed into blood stream via rumen wall and converted to glucose in the liver. Glucose formed is used by animal as energy source for maintenance, production and reproduction activities (McDonald et al., 1987). In addition, Weinberg et al. (2004) concluded that LAB from silage has potential role as probiotic which beneficially affects the host animal by improving its intestinal microbial balance.

Concentration of lactic acid in silage treated with combination of LAB and acacia tannin (C, D, E and F) tended to be lower in comparison with silage treated with LAB alone (B). McSweeney et al. (2001) revealed that tannin has ability to bind to macromolecules such as structural carbohydrate and starch, thereby impairing their degradation. However, concentration of lactic acid in king grass silages treated LAB and acacia tannin were slightly lower than the ideal range of lactic acid concentration from 80 to 120 g/kg DM.

Concentration of NH₃-N in silages treated with LAB or combined with acacia tannin (B, C, D, E and F) were lower (P<0.05) than that of control silage. The finding was consistent with previous work of Nishino and Uchida (1999) who found that use of fermented Lucerne extract increased lactic acid and greatly inhibited the clostridial activity to protect proteins from extensive degradation. As stated by Ohshima and McDonald (1978) that during ensiling protein is degraded to peptides and free amino acids by plant proteases. In addition, degradation of amino acids to ammonia and non-protein nitrogenous fraction is predominantly due to proteolytic clostridia. Chamberlain and Wilkinson (1996) concluded that ammonia-N is as an indicator of the proportion of the total N which has been completely degraded during ensiling. Hence, concentration of ammonia-N is the best indicator of secondary fermentation. The growth of proteolytic clostridia, which degrade protein and amino acids to NH₃, is inhibited by low pH (Ohshima and McDonald, 1978; McDonald et al., 1987). This result was supported by the low pH values in silage B, C, D, E and F compared to silage A. This condition

could depress the growth of proteolytic clostridia. Concentration of NH₃-N reduced with increasing tannin addition in silage materials. This could be due to tannin can bind to protein and protect them from microbial degradation. Salawu et al. (1999) reported that tannin were able to protect herbage proteins from plant/microbial enzyme hydrolysis during ensiling. Concentration of NH₃-N in silage C, D, E and F was decreased with increasing tannin concentration added in silage. This result is in agreement with previous study by Tabacco et al. (2006) that concentrations of NH₃-N and NPN were reduced with increasing tannin concentration in silage. The normal range of NH₃-N concentration in silage is 50 to 150 g NH₃-N/kg DM. However, the target value for NH₃-N is less than 50 g/kg total N (Chamberlain and Wilkinson, 1996). Based on NH₃-N concentration, silage C, D, E and F could be classified in normal range of NH₃-N concentration.

The VFAs comprise of acetic acid, propionic acid, butyric acid and other acids. The production of these acids is a reflection of an inefficient fermentation or of secondary fermentation of lactic acid to butyric acid and degradation of amino acids to ammonia with the production of acetic acid from the carbon skeleton of the amino acid (Chamberlain and Wilkinson, 1996). Concentrations of total VFA, acetic acid and propionic acid were not significantly different among treatments silage. However, silage treated with epiphytic LAB and tannin had lower ($P<0.01$) butyric acid concentration compared to control silage. The reduced butyric acid in those silages could be due to lower pH silage which may inhibit the activity of clostridia. McDonald (1991) stated that reducing pH silage prevented the growth of undesirable microbes *e.g.* listeria, clostridia, enterobacteriaceae and moulds. Another possible explanation for the reduction of butyric acid in silage treated with acacia tannin (silage C, D, E and F) might be due to inhibited the activity of clostridia. It is well known that clostridia are responsible for most of the butyric acid in silage (McDonald, 1981).

Visually, there were no moulds on the top of silages treated with acacia tannin. This was probably due to toxic effect of tannin to silage moulds. This result is supported with previous study of Salawu et al. (1999) that higher tannin in the silage reduced the activity of moulds.

Chemical Composition of Silages

Chemical composition of silages treated with epiphytic LAB and tannin of acacia is given at Table 3.

Dry matter content in six silages was still lower than the target value for DM in ideal silage as recommended by Chamberlain and Wilkinson (1996). The lower DM content in all silages could be attributed to DM content of king grass used as silage material was less than 20%. The OM content in silage A was similar to silages treated with epiphytic LAB and tannin of acacia. Concentration of CP in silages treated with epiphytic LAB and tannin of acacia was higher ($P<0.01$) than that of control silage. The higher CP content in those silages could be due to lower activity proteolytic as resulted by low in pH value and protein protection by tannin acacia. Other explanation for higher CP content in silage treated with epiphytic LAB and tannin of acacia is addition of protein obtained from grass and acacia extracts.

Table 3. Characteristics of silages after ensiling for 30 days

	Silages						SE	P
	A	B	C	D	E	F		
DM	17.5 ^{ab}	17.5 ^{ab}	18.2 ^a	16.2 ^d	16.8 ^{bc}	17.9 ^a	0.32	0.02
OM	94.6	95.3	95.5	96.2	95.2	95.4	0.45	0.27
CP	12.8 ^B	16.2 ^A	15.7 ^A	15.3 ^A	15.7 ^A	15.7 ^A	0.19	<0.01
NDF	71.0 ^A	65.3 ^B	68.0 ^{AB}	67.3 ^B	67.9 ^{AB}	66.9 ^B	0.96	<0.01
ADF	43.4 ^a	41.7 ^{abc}	40.2 ^c	40.7 ^{bc}	42.5 ^{ab}	41.8 ^{abc}	0.55	0.02
Hemicellulose	28.6 ^A	23.5 ^B	27.8 ^A	26.6 ^{AB}	25.4 ^{AB}	25.1 ^{AB}	0.79	<0.01

^{a-b} Means in the same row followed by different letters are different ($P<0.05$)

^{A-B} Means in the same row followed by different letters are different ($P<0.01$)

The NDF content in silages B, D and F was lower ($P<0.01$) than that of control silage. Silages C and D had lower ($P<0.05$) ADF content compared with control silage. This results, however is in agreement with previous study of Yahaya et al. (2004b). One of the explanations for the lower NDF

and ADF in those silage is that enzymatic action *e.g.* hemicellulases, cellulase present in the original forage on cell wall during ensiling. Decreased NDF and ADF concentrations in silage treated with epiphytic LAB or combined with tannin of acacia had beneficial effect of silage nutritive value and leading to an increase in silage digestibility in the rumen.

CONCLUSIONS

Fermented grass extract prepared from king grass could be used as a good source of lactic acid bacteria pre-ensiling. Additions of epiphytic LAB or combined with acacia tannin have beneficial effect on fermentation and nutritive quality of king grass silage. This was indicated by a higher lactic acid production and lower pH value, as well as concentrations of NH₃-N and butyric acid. Silages treated with epiphytic LAB or combined with tannin of acacia had a higher crude protein content and lower cell wall component contents *i.e.* NDF, ADF and hemicellulose compared with control silage.

LITERATURE CITED

- AOAC. 1990. Official Methods of Analysis of the Association of Analytical Chemist. 16th ed. Association of Official Analytical Chemist, Arlington, VA, USA.
- Barker, S.B. and W.H. Summerson. 1941. The colorimetric determination of lactic acid in biology material. *J. Biol. Chem.* 138: 535-554.
- Bureenok, S., T. Namihira, S. Mizumachi, Y. Kawamoto and T. Nakada. 2006. The effect of epiphytic lactic acid bacteria with or without different byproduct from defatted rice bran and green tea waste on napiergrass (*Pennisetum purpureum* Shumach) silage fermentation. *J. Sci. Food Agric.* 86: 1073-1077.
- Chamberlain, A.T. and J.M. Wilkinson. 1996. *Feeding the Dairy Cow*. Chalcombe Publications, Lincoln, UK.
- Chamberlain, D.G. 1987. The silage fermentation in relation to the utilization of the nutrients in the rumen. *Process Biochem.* 22: 60-63.
- Chaney, A.L. and E.P. Marbach. 1962. Modified reagents for determination of urea and ammonia. *Clin. Chem.* 8: 130-132.
- Hariadi, B. Tj. and B. Santoso. 2010. Evaluation of tropical plants containing tannin on in vitro methanogenesis and fermentation parameters using rumen fluid. *J. Sci. Food Agric.* 90:456-461.
- Harrison, J. H. and R. Blauwiekel. 1994. Fermentation and utilization of grass silage. *J. Dairy Sci.* 77:3209-3235.
- Kondo, M., K. Kita and H. Yokota. 2004. Feeding value to goats of whole-crop oat ensiled with green tea waste. *Anim. Feed Sci. Technol.* 113: 71-81.
- Lin, K., K. Bolsen, B.E. Brent, R.A. Hart and J.T. Dickerson. 1992. Epiphytic microflora on alfalfa and whole-plant corn. *J. Dairy Sci.* 75: 2484-2493.
- Masuko, T., Y. Hariyama, Y. Takahashi, L. M. Cao, M. Goto and M. Ohshima. 2002. Effect of addition of fermented juice of epiphytic lactic acid bacteria prepared from timothy and orchardgrass on fermentation quality of silages. *Grassl. Sci.* 48 (2): 120-125.
- McDonald, P., A.R. Hunderson and S.J.E. Heron. 1991. *The Biochemistry of Silage*. 2nd ed. Cambrian Printers Lt. Aberystwyth. Pp. 184-236.
- McDonald, P., R.A. Edwards and J.F.D. Greenhalgh. 1987. *Animal Nutrition*. 4th ed. Longman Group Ltd. Hongkong.
- McSweeney, C.S., B. Palmer, D.M. McNeill and D.O. Krause. 2001. Microbial interactions with tannins: nutritional consequences for ruminants. *Anim. Feed Sci. Technol.* 91: 83-93.
- Muck, R.E. 1990. Prediction of lactic acid bacteria numbers on lucerne. *Grass Forage Sci.* 45: 273-280.
- Nishino, N. and S. Uchida. 1999. Laboratory evaluation of previously fermented juice as a fermentation stimulant for lucerne silage. *J. Sci. Food Agric.* 79: 1285-1288.
- Ohshima, M. and P. McDonald. 1978. A review of the changes in nitrogenous compounds of herbage during ensilage. *J. Sci. Food Agric.* 29: 497-505.
- Ohshima, M., E. Kimura and H. Yokota. 1997. A methods of making good quality silage from direct cut alfalfa by spraying previously fermented juices. *Anim. Feed Sci. Technol.* 66: 129-137.
- Salawu, M.B., T. Acamovic, C.S. Stewart, T. Hvelpund and M.R. Weisbjerg. 1999. The use of tannins as silage additives: effects on silage composition and mobile bag disappearance of dry matter and protein. *Anim. Feed Sci. Technol.* 82: 243-259.
- Santoso, B. B. Tj. Hariadi, H. Manik and H. Abubakar. 2009. Quality of tropical grasses ensiled with additive of lactic acid bacteria prepared from fermented grass extract. *J. Anim. Sci. Technol.* 32(2): 138-145. (in Indonesian)
- Santoso, B., M N. Lekitoo and Umiyati. 2007. Chemical composition and nutrient degradation of elephant grass

- silage ensiled with black tea leaf residue. *Anim. Prod.* 9(3): 160-165. (in Indonesian)
- Seglar, B. 2003. Fermentation Analysis and Silage Quality Testing Proceeding of the Minnesota Dairy Health Conference College of Veterinary Medicine, University of Minnesota. pp. 119-136.
- Smith, L.H. 1962. Theoretical carbohydrate requirement for alfalfa silage production. *Agron. J.* 54 (4): 591-593.
- Tabbaco, E., G. Borreani, G.M. Crovetto, G. Galassi, D. Colombo and L. Cavallarin. 2006. Effect of chestnut tannin on fermentation quality, proteolysis, and protein rumen degradability of alfalfa silage. *J. Dairy Sci.* 89: 4736-4746.
- Tamminga, S. 1992. Nutrition management of dairy cows as a contribution to pollution control. *J. Dairy Sci.* 75: 345-357.
- Van Soest, P.J., J.B. Robertson and B.A. Lewis. 1991. Methods for dietary fiber, neutral detergent fiber, and nonstarch polysaccharides in relation to animal nutrition. *J. Dairy Sci.* 74: 3583-3597.
- Wang, J., J.Q. Wang, H. Zhou and T. Feng. 2009. Effects of addition of previously fermented juice prepared from alfalfa on fermentation quality and protein degradation of alfalfa silage. *Anim. Feed Sci. Technol.* 151: 280-290.
- Weinberg, Z.G., R.E. Muck, P.J. Weimer, Y. Chen and M. Gamburg. 2004. Lactic acid bacteria used in inoculants for silage as a probiotics for ruminants. *Appl. Biochem. Biotech.* 118: 1-10.
- Yahaya, M.S., M. Goto, W. Yimiti, B. Smerjai and Y. Kuwamoto. 2004a. Additive effects of fermented juice of epiphytic lactic acid bacteria and acetic acid on silo fermentation and ruminal degradability of tropical elephant grass. *J. Anim. Vet. Adv.* 3: 116-122.
- Yahaya, M.S., M. Goto, W. Yimiti, B. Smerjai and Y. Kuwamoto. 2004b. Evaluation of fermentation quality of a tropical and temperate forage crops ensiled with additives of fermented juice of epiphytic lactic acid bacteria (FJLB). *Asian-Aust. J. Anim. Sci.* 17: 942-946.